


RESEARCH

Open Access



Fear and depression during the COVID-19 outbreak in Cameroon: a nation-wide observational study

Joseph Nelson Siewe Fodjo^{1,2}, Leonard Ngarka², Wepnyu Y. Njamnshi^{2,3}, Leonard N. Nfor², Michel K. Mengnjo², Edwige Laure Mendo², Samuel A. Angwafor², Jonas Guy Atchou Basseguin², Cyrille Nkouonlack², Edith N. Njit², Nene Ahidjo², Eric S. Chokote², Fidèle Dema², Julius Y. Fonsah², Godwin Y. Tatah², Nancy Palmer², Paul F. Seke Etet², Dennis Palmer², Dickson S. Nsagha², Daniel E. Etya'ale³, Stephen Perrig³, Roman Sztajzel³, Jean-Marie Annoni³, Anne-Cécile Zoung-Kanyi Bissek⁴, Rose G. F. Leke², Marie-Thérèse Abena Ondo Obama², John N. Nkengasong⁵, Robert Colebunders¹ and Alfred K. Njamnshi^{2,3*} 

Abstract

Background: The COVID-19 pandemic has been associated with significant psychological and social distress worldwide. We investigated fear and depression among adults in Cameroon during different phases of the COVID-19 outbreak.

Methods: An online survey was conducted in Cameroon from June–December 2020 using a structured questionnaire. Socio-demographic data and information regarding COVID-19 history were obtained. Fear and depressive symptoms were assessed using the Fear of COVID-19 score (FCV-19S) and the Patient Health Questionnaire (PHQ-9), respectively. Responses were clustered in weeks to better appreciate their evolution over time.

Results: Overall, 7381 responses from all ten regions of Cameroon were analysed (median age: 30 years, 73.3% male). The prevalence of depression (PHQ-9 score ≥ 10) was 8.4%, and that of high fear of COVID-19 (FCV-19S scores ≥ 19) was 57.4%. These rates were similar across genders, age-groups, and region of residence. While mean weekly PHQ-9 scores remained fairly stable throughout the study period (range: 2.53–3.21; $p = 0.101$), mean FCV-19S scores were highest during the early weeks but decreased significantly thereafter (from 20.31 to 18.34; $p < 0.001$). Multivariate analyses revealed that having a postgraduate degree, a history of quarantine, flu-like symptoms during the past 14 days, and higher FCV-19S scores were associated with more severe depressive symptoms, while obtaining COVID-19 information from various sources reduced the odds for depression.

Conclusion: Depression amidst the COVID-19 crisis is less prevalent in Cameroon than in other countries. Prompt and widespread dissemination of adequate COVID-19 information may reduce the risks for depression by dispelling fear and anxiety among Cameroonians.

Keywords: COVID-19, PHQ-9, FCV-19S, Cameroon, Fear, Depression

* Correspondence: alfred.njamnshi@brainafrica.org

²Brain Research Africa Initiative (BRAIN), Yaoundé, Cameroon

³Brain Research Africa Initiative (BRAIN), Geneva, Switzerland

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Since its initial outbreak in China in December 2019, the coronavirus disease 2019 (COVID-19) has spread globally and caused over a million deaths [1]. Unprecedented restrictive measures were deployed by every nation to prevent or contain local epidemics. Besides the biological (and sometimes fatal) damages caused by the virus, the COVID-19 pandemic has been associated with significant psychological and social distress around the globe. Indeed, studies conducted during the COVID-19 pandemic have revealed high levels of stress, anxiety and depressive disorders among the general population [2], as well as specific populations: healthcare workers and their families [3–5], military personnel [6], and COVID-19 survivors [7]. Available data suggests that women and younger individuals are more affected by psychosocial symptoms during the COVID-19 pandemic [8, 9]. The socio-economic consequences of the strict lockdown measures implemented in many countries further exacerbated the financial insecurities of the population with adverse effects on their psychological and social well-being, leading to suicide in some cases [10].

Cameroon, a Low-Middle income country located in the Central African sub-region, was not spared by the COVID-19 pandemic. After reporting its first case on March 6th 2020, community transmission was observed resulting in a cumulative 26,277 cases and 448 deaths as of December 31st 2020 [11]. The necessary preventive measures prescribed by the Cameroonian government [12] have disrupted daily routines and adversely impacted the nation's socio-economic landscape given that a large proportion of the population relies on the informal sector for its livelihood. Although the mental or brain health impacts of the COVID-19 pandemic are evident in Cameroon [13], the magnitude of the problem is still poorly understood as no nation-wide study has yet been done. We therefore conducted this study to assess the level of fear and the frequency of depressive disorders in the Cameroonian population amidst the COVID-19 outbreak. We further sought to identify determinants of fear and depressive symptoms during the ongoing health crisis.

Methods

Study setting and population

The study was conducted from June 5th to December 5th 2020, and recruited participants from all ten regions of Cameroon. Cameroon's population has a median age of 18.7 years and life expectancy of 60.3 years, with 56.3% of people living in urban settings [14]. For over 4 years, the nation has been confronted with an internal conflict in the two English-speaking regions (North West and South West) and sporadic terrorist attacks in the Northern part of the country.

Study tools and procedures

We performed online surveys designed by Brain Research Africa Initiative (BRAIN) researchers in collaboration with the International Citizen Project on COVID-19 (ICP-Covid) consortium [15]. The ICPCovid website, initiated by a team of researchers based at the University of Antwerp in Belgium, offers a secure electronic platform to collect COVID-19-related data from several low- and middle-income countries in order to assess the population's perception of, and adherence to the implemented preventive measures. A web-based online questionnaire was designed using the 'Drupal' system, translated to English and French and pre-tested on Cameroonian adults. Besides collecting the socio-demographic information of participants, we also assessed their psychological and social well-being using two scales:

- Fear of COVID-19 scale (FCV-19S): This is a 7-item tool that was recently validated in Iran [16]. Each item on this scale is scored between 1 and 5. A total score is calculated by adding up each item score (overall score ranging from 7 to 35). The higher the score, the greater the fear of coronavirus-19. A cut-off score of 19 and above on this scale has been suggested to identify individuals with a high level of COVID-19-related fear [17].
- Patient Health Questionnaire-9 (PHQ-9): This 9-item tool was developed to screen for major depressive disorders among adults [18]. Each of the nine items is scored between 0 and 3 (maximum possible score: 27). Scores of 10 or above are often considered as an indication of a likely depressive disorder, as previously validated in a Cameroonian study population [19]. Other cut-offs have been proposed to further discriminate the severity of depressive symptoms: cut-off scores of 0–4 = none/minimal; 5–9 = mild; 10–14 = moderate; 15–19 = moderately severe; 20–27 = severe [3, 17].

In addition to the above-mentioned screening tools, Likert-format questions were asked regarding the level of worry experienced by the respondents regarding their own health and the health of their loved ones during the COVID-19 crisis; this was expressed on a 5-point scale ranging from 1 (not at all worried) to 5 (extremely worried). The web-link to the electronic survey was disseminated via social media platforms and bulk messaging to phone users. Upon clicking on the link, the user was directed to an information and consent page where he/she could agree to participate, fill in the responses submitted via a smartphone, tablet, or computer. The electronic questionnaire was made accessible during certain periods each month, and closed down intermittently between survey rounds. All submitted responses were

immediately stored in a password-protected server in Belgium until data retrieval.

Data analysis

Collected data were exported to Microsoft Excel 2016 spreadsheets for cleaning, and later transferred to R version 4.0.2 for analysis. Based on the fact that continuous variables were not normally distributed as shown by the Kolgomorov-Smirnov test, we summarized them as median with interquartile range (IQR). On the other hand, categorical variables were expressed as fractions and percentages. To investigate determinants of depressive symptoms as screened by the PHQ-9 tool, an ordinal logistic regression model was constructed with the different severity levels of depression (none/minimal, mild, moderate, moderately severe, severe) as dependent variable and socio-demographic variables as covariates. To capture the effect of the ongoing armed conflicts in some parts of the country, a binary variable was created as follows: conflict-stricken region = yes (for respondents residing in the Far North, North West, and South West regions) vs no (for all other regions). Given the prolonged study period and the rapid evolution of COVID-19 dynamics, we took into account the timing of the responses by considering the different study weeks as clusters. Only weeks with ≥ 100 responses were included in the multivariate analysis. We used the *polr* function (package: 'MASS') and the *vcovCL* function (package: 'sandwich') in the software R to obtain clustered standard errors from the regression model. Covariates for the final model were selected based on a p -value < 0.2 during univariate analysis.

Results

Of the 7538 responses received, 7381 were eligible for analysis (median age: 30 years, 73.3% male); Table 1. About one-fifth of the participants ($n = 1454$; 19.7%) resided in the conflict-stricken regions of Cameroon. Majority of participants (70.9%) had attained university level of education.

The median PHQ-9 score in our study population was 1.0 (IQR: 0–4) on a scale ranging from 0 to 21. Using the PHQ-9 cut-off value of ≥ 10 as positive screening for depression, the prevalence of depression was 617/7381 (8.4%). Applying the other cut-offs revealed the frequencies of no/minimal, mild, moderate, moderately severe, and severe depression to be respectively: 5669 (76.8%), 1095 (14.8%), 346 (4.7%), 178 (2.4%), and 93 (1.3%). Regarding COVID-19 fear, FCV-19S scores ranged from 7 to 35, with median 20 (IQR: 15–23) and mean 19.3 ± 6.4 . Furthermore, more than half of the respondents ($n = 4238$; 57.4%) reported experiencing high levels of fear (i.e. FCV-19S scores ≥ 19). The prevalence of both depression (PHQ-9 score ≥ 10) and high level of fear (FCV-19S ≥ 19) were similar across genders, age groups,

Table 1 Participants' characteristics

Characteristics	Survey findings N = 7381
Age: Median [IQR]	30.0 (25.0–38.0)
Gender: n (%)	
Male	5409 (73.3%)
Female	1972 (26.7%)
Highest educational level: n (%)	
Primary school	118 (1.6%)
Secondary school	2029 (27.5%)
University: Undergraduate	2963 (40.1%)
University: Postgraduate	2271 (30.8%)
Residential setting: n (%)	
Rural	672 (9.1%)
Sub-Urban	1327 (18.0%)
Urban	5382 (72.9%)
Living alone in household: n (%)	1383 (18.7%)
Self-reported socio-economic status: n (%)	
Low class	2494 (33.8%)
Lower-middle class	3839 (52.0%)
Upper-middle class	912 (12.4%)
High class	136 (1.8%)
Profession: n (%)	
Student	1832 (24.8%)
Unemployed	1342 (18.2%)
Self-employed	869 (11.8%)
Private employee	1843 (25.0%)
Government employee	1285 (17.4%)
Retired	210 (2.9%)
Healthcare worker or student: n (%)	793 (10.7%)
Source of COVID-19 information: n (%) ^a	
Radio, Television, or government announcements	6601 (89.4%)
Social Media	5780 (78.3%)
Healthcare worker	2874 (38.9%)
Underlying chronic disease: n (%) ^b	716 (9.7%)
History of being quarantined at home/institution: n (%)	1375 (18.6%)
History of violence/discrimination during COVID-19 outbreak: n (%)	1334 (18.1%)

IQR Interquartile range [25th percentile – 75th percentile]

^aEach participant was allowed to choose more than one answer, hence the categories may overlap

^bHeart disease, diabetes, hypertension, cancer, HIV, or asthma

residential setting, being in the healthcare sector or not, and region of residence (Table 2).

FCV-19S scores correlated significantly with PHQ-9 scores (Spearman-rho = 0.28, $p < 0.001$). Considering only weeks with at least 100 responses, both COVID-19

Table 2 Prevalence of depression and high fear levels during the COVID-19 outbreak in Cameroon

Characteristics		FCV-19S ≥ 19 : n (%)	P-value ^a	PHQ-9 score ≥ 10 : n (%)	P-value ^a
Gender	Male	3079 (56.9%)	0.163	458 (8.5%)	0.611
	Female	1159 (58.8%)		159 (8.1%)	
Age	18–25 years	1186 (56.7%)	0.289	185 (8.8%)	0.588
	26–35 years	1724 (57.4%)		254 (8.5%)	
	36–45 years	750 (56.4%)		104 (7.8%)	
	46–55 years	365 (60.9%)		51 (8.5%)	
	> 55 years	213 (60.0%)		23 (6.5%)	
Region	Centre	1267 (57.8%)	0.758	178 (8.1%)	0.773
	Adamawa	183 (57.4%)		29 (9.1%)	
	East	101 (55.8%)		9 (5.0%)	
	Far North	247 (58.1%)		43 (10.1%)	
	Littoral	1006 (57.0%)		147 (8.3%)	
	North	131 (55.3%)		19 (8.0%)	
	North West	414 (56.6%)		67 (9.2%)	
	South	129 (54.9%)		18 (7.7%)	
	South West	440 (60.9%)		59 (8.2%)	
	West	320 (55.8%)		48 (8.4%)	
Residential setting	Rural	374 (55.7%)	0.624	51 (7.6%)	0.646
	Sub-urban	765 (57.6%)		107 (8.1%)	
	Urban	3099 (57.6%)		459 (8.5%)	
Worker/student in healthcare	Yes	432 (54.5%)	0.083	67 (8.5%)	0.977
	No	3806 (57.8%)		550 (8.4%)	

^aChi-Squared test

fear and PHQ-9 scores seemed to increase with increasing weekly incidence of COVID-19 in Cameroon, and vice versa (Fig. 1). While the COVID-19 weekly incidence in Cameroon did not correlate significantly with the respondents' FCV-19S scores (Spearman-rho = 0.021, $p = 0.078$), a weak positive correlation was found between weekly COVID-19 incidence and PHQ-9 scores (Spearman-rho = 0.033, $p = 0.006$). Mean weekly PHQ-9 scores did not vary significantly across the study period (range: 2.53–3.21; $p = 0.101$), in contrast to mean FCV-19S scores which were highest during the early weeks of the survey but decreased significantly thereafter (range: 18.34–20.31; $p < 0.001$). The median Likert score for worrying about other people's health was significantly higher than that of worrying for one's own health: 1 (IQR: 1–3) vs 1 (IQR: 1–2), $p < 0.001$.

Using dichotomized PHQ-9 and FCV-19S scores, we estimated the prevalence of depression and high fear for the study weeks with ≥ 100 responses (Table 3). We observed similar proportions of respondents with depression across the different study weeks ($p = 0.195$), while there were significant differences in the prevalence of high fear ($p < 0.001$).

The multivariate analysis revealed that having a post-graduate degree, a history of quarantine, experiencing flu-like symptoms during the past 14 days, and higher FCV-19S scores were associated with increased odds to have more severe depressive symptoms. Meanwhile, obtaining COVID-19 information from either official sources or from the social media reduced these odds (Table 4).

Discussion

Our study is the first to assess fear and depression among Cameroonian adults during 6 months of the COVID-19 crisis, in a nation-wide large survey. Overall, almost one-tenth of respondents screened positive for depression and the PHQ-9 depression scores appeared to vary proportionately to the weekly incidence of COVID-19. It is worth noting that during the latter months of the study period, schools were allowed to resume cautiously and this brought about a semblance of normalcy in the daily routines of Cameroonians. These changes may have had repercussions on the overall psychosocial well-being of the study participants over time.

The prevalence of depression during the COVID-19 crisis in our study is lower compared to previous reports




Fig. 1 Weekly evolution of COVID-19 incidence and: **A** Fear scores; **B** PHQ-9 scores

by researchers who also used the PHQ-9 tool with a cut-off score of ≥ 10 in both high-income and low- middle-income countries [17, 20–22]. Our numbers are also lower than pre-COVID-19 depression rates in healthy Cameroonian adults, estimated at 19.8% in one study [23] but the fact that a different screening tool (Beck's Depression Inventory) was used renders comparisons with our study difficult. It is expected that the COVID-19

pandemic would cause a rise in the burden of depression compared to baseline levels, as observed during a comparative study in American adults [24]. Of note, pre-COVID-19 depression levels were already low in Africa compared to other continents (except for Australia) [25], and the pandemic would likely exacerbate existing situations without necessarily altering the global distribution of depressive disorders. Furthermore, in resource-limited settings such as Cameroon where economic hardship is frequent, lower depression rates may also be observed because people are more focused on being financially productive amidst the crisis to keep providing for their families, even to the point of overlooking or mentally downplaying the threat of COVID-19. This notwithstanding, the most educated respondents (post-graduate level) who probably had a better understanding of the COVID-19 science and the risks associated with the pandemic were prone to more severe depressive symptoms (adjusted odds ratio of 1.5, relative to respondents with primary level education).

Regarding FCV-19S results, over half of the participants experienced high levels of fear vis-à-vis COVID-19. This is greater than the 35.7% reported in Greece [17] using a similar methodology. COVID-19 fear scores were highest at the beginning of the study period (June 2020), possibly because that period represented the first epidemic peak in Cameroon with thousands of cases confirmed on a weekly basis. No regional disparity was noted in the prevalence of depression or high fear of COVID-19. This is an intriguing finding, because we expected that persons residing in conflict-stricken regions would experience more psychological and social distress in the face of the COVID-19 outbreak. Qualitative research may be required to better understand the determinants of mental and brain health, as well as the coping mechanisms of individuals in different settings during the COVID-19 pandemic.

Table 3 Prevalence of depression and high fear across the study weeks

Calendar week	High fear (FCV-19S ≥ 19): n (%)	P-value ^a	Depression (PHQ-9 score ≥ 10): n (%)	P-value ^a
Week 30 (n = 817)	517 (63.3%)	< 0.001	74 (9.1%)	0.195
Week 31 (n = 286)	167 (58.4%)		27 (9.4%)	
Week 34 (n = 1132)	677 (59.8%)		84 (7.4%)	
Week 35 (n = 120)	68 (56.7%)		11 (9.2%)	
Week 36 (n = 640)	333 (52.0%)		32 (5.0%)	
Week 39 (n = 152)	90 (59.2%)		14 (9.2%)	
Week 40 (n = 454)	248 (54.6%)		36 (7.9%)	
Week 43 (n = 1686)	940 (55.8%)		159 (9.4%)	
Week 47 (n = 1347)	785 (58.3%)		132 (9.8%)	
Week 49 (n = 397)	216 (54.4%)		26 (6.6%)	

^aChi squared test

Table 4 Clustered ordinal logistic regression investigating factors associated with increasing severity of depressive symptoms

Covariates	Univariate <i>P</i> -value	Adjusted OR (95% CI)	Multivariate <i>P</i> -value
Age (in years)	0.099	0.994 (0.986–1.003)	0.177
Socioeconomic status:			
Low	Ref	Ref	
Lower middle	0.097	1.137 (0.991–1.305)	0.066
Upper middle	0.809	1.041 (0.853–1.269)	0.694
High	0.117	1.300 (0.853–1.980)	0.222
Educational level			
Primary	Ref	Ref	
Secondary	0.337	1.354 (0.914–2.006)	0.131
Undergraduate	0.247	1.388 (0.919–2.097)	0.119
Postgraduate	0.141	1.523 (1.011–2.294)	0.044
Profession			
Student	Ref	Ref	
Jobless	0.767	1.026 (0.877–1.201)	0.746
Self-employed	0.173	0.974 (0.787–1.205)	0.808
Private worker	0.850	1.022 (0.928–1.127)	0.655
Government worker	0.614	0.953 (0.819–1.107)	0.526
Retired	0.914	1.210 (0.727–2.013)	0.463
Student / worker in healthcare sector	0.159	1.095 (0.958–1.252)	0.185
COVID-19 information from official sources (TV, radio, or other government channels)	0.011	0.801 (0.728–0.882)	< 0.001
COVID-19 information from social media	0.078	0.831 (0.758–0.912)	< 0.001
Living alone in household	0.132	1.098 (0.986–1.224)	0.089
Flu-like symptoms during the past 14 days	< 0.001	2.296 (2.050–2.570)	< 0.001
History of quarantine/isolation for COVID-19	< 0.001	1.568 (1.422–1.728)	< 0.001
History of violence/discrimination	0.182	0.949 (0.816–1.104)	0.501
Fear of COVID-19 score	< 0.001	1.114 (1.102–1.127)	< 0.001
Residential setting		–	–
Rural	Ref		
Sub-urban	0.639		
Urban	0.567		
COVID-19 information from healthcare workers	0.929	–	–
Male gender	0.459	–	–
Conflict-stricken region	0.329	–	–

OR Odd's ratio, CI Confidence interval, Ref Reference category

The frequency of depression and high fear of COVID-19 was similar among participants who were workers/students in the healthcare sector and others not involved in healthcare. A possible explanation for this observation is the reduced exposure to COVID-19 patients for health personnel in Cameroon due to a relatively lower COVID-19 burden compared to other regions of the world. Indeed, increased exposure to COVID-19 cases has been shown to fuel stress, anxiety and depression among healthcare providers [26]. Therefore, it is

understandable that mental and brain disorders during the COVID-19 pandemic may be more frequent among healthcare workers in Europe, America, Asia [3, 6, 27–29], but also in neighbouring Nigeria [30] which has more than thrice the number of COVID-19 cases in Cameroon [1]. It is worth noting that another survey previously conducted among healthcare workers in Cameroon during the COVID-19 pandemic found a prevalence of depression of 42.8% using the Hospital Anxiety and Depression scale (HADS) [31], compared to

8.5% in our study. While acknowledging the methodological differences between our study and the aforementioned survey, their findings do suggest that exposure of healthcare workers to COVID-19 patients is an important stressor in Cameroon. The authors reported that fear of contamination with the coronavirus and fear of death were modulators of depression among the health personnel [31].

Participants with a history of quarantine, or those who had recently experienced flu-like symptoms (during the past 2 weeks) were more likely to exhibit depressive symptoms. These results concur with observations made in Canada during the SARS-1 outbreak during which longer quarantine durations and a history of contact with an infected person were associated with increased psychological distress [32]. Similar observations were made in Brazil and Portugal during the COVID-19 pandemic [33]. This highlights quarantine / isolation as a risk factor for short- and possibly long-term psychosocial distress, requiring careful consideration by the health authorities [34]. We observed that during the COVID-19 outbreak in Cameroon, the respondents were more concerned about the health of their loved ones than their own health. This finding is relevant in the context of implementing public health measures such as mandatory quarantine and vaccination for COVID-19. Although adherence to preventive measures may depend more on their practicality (ease of implementation) than on the psychology of individuals [35], one could still appeal to altruism by reminding the public about the community-wide benefits of such interventions. It is clear the impact of the interventions goes beyond individual protection to shielding other community members from becoming infected. In a situation whereby even healthcare workers are against a vaccine trial in their community [36], it is important to emphasize the collective benefits of vaccination while rendering the process of COVID-19 vaccination as simple and attractive as possible in a bid to improve acceptability. Studies on attitudes and acceptability of the COVID-19 vaccine in Cameroon are needed.

Participants who resorted to various platforms (radio, television, government announcements, or social media) in search of information about COVID-19 had reduced odds of experiencing depressive symptoms. Similar findings were reported by González-Sanguino et al. in Spain, as they found a negative relationship between depression and having adequate information about the coronavirus [9]. Meanwhile, a study conducted among Chinese adults during their nationwide lockdown showed that a higher level of COVID-19 media exposure was significantly associated with higher PHQ-9 scores [37]. We surmise that the information from various media rather produced a soothing effect on Cameroonian participants

by unveiling how grave the COVID-19 situation was in other countries, and providing a sense of relative safety. Indeed, the COVID-19 burden and death toll in Cameroon is considerably lower compared to the daily reports from Europe and the Americas [38]. It is expedient that the media outlets be leveraged to promptly disseminate adequate COVID-19-related information to the public, as this approach is a pillar in ensuring optimal mental and brain health during such health crises by addressing any sense of uncertainty and fear [39].

A number of limitations must be taken into account when interpreting our study findings. This was an online survey, and it is therefore impossible to verify the veracity of the provided responses. In addition, we acknowledge a sampling bias since our survey could only be filled by literate and educated individuals who had access to internet. Indeed, the web-based recruitment procedures could not achieve a representative study population. Fewer female respondents participated in the survey; this could be due to the fact that in Cameroon, it was reported that women are less likely to use the internet compared to men [40]. After the preliminary data analysis, an effort was made through the Ministry of Women's Empowerment and the Family to encourage more women in women's groups to participate but the percentage increase in participation was just 1.5 (from 25.2 to 26.7). Since the female gender has been associated with greater psychological distress during the COVID-19 outbreak [9], the male predominance in our study population might have led to an underestimation of the overall burden of COVID-19-related fear and depression in Cameroon. Lastly, the survey's duration was very long, considering the rapidly evolving dynamics of COVID-19 and could lead to participant fatigue. However, this was mitigated by constituting weekly clusters during the data analysis.

Conclusions

Our study found that less than 10% of Cameroonian adults reported depressive symptoms amidst the COVID-19 crisis, much lower than what has been documented in other settings. Meanwhile, over half of the respondents experienced a high level of fear of COVID-19. We recommend that various media outlets should be leveraged to disseminate adequate information about COVID-19 as this will dispel fear and reduce the risk for depression. Furthermore, persons who report flu-like symptoms, the more educated individuals (postgraduates) and those who had been quarantined seem to be at increased risk for depression, and should be prioritized for interventions during the COVID-19 pandemic in Cameroon. Finally, appealing to the population's sense of altruism may be the way forward to ensure optimal

mental and brain health, as well as better acceptability of large scale COVID-19 preventive measures.

Abbreviations

COVID-19: Coronavirus disease 2019; FCV-19S: Fear of COVID-19 score; ICPCovid: International Citizen Project on COVID-19; IQR: Interquartile range; PHQ-9: Patient Health Questionnaire

Acknowledgements

We are grateful to the Cameroon Prime Minister's Office for overall administrative support, the Ministry of Communication and the Ministry of Post and Telecommunications of Cameroon and its partners for assisting in sensitisation of the population and in disseminating the survey link. We appreciate the Ministry of Women's Empowerment and the Family for sensitisation through women's organisations. We sincerely thank the participants for providing the responses.

Authors' contributions

SFJN, CR, and NAK conceived the study. NAK obtained administrative support/ethical clearance in Cameroon and supervised the entire project. NL, NWY, LNN MMK and MEL coordinated data collection. ASA, ABJG, NC, NEN, AN, CES, DF, FJY, TGY, PN, SEPF, PD, NDS, EDE, PS, SR, AJM, Z-KBAC, LRGF, AOOMT & NNJ contributed to adaptation of research instrument to the Cameroonian context and coordinated sensitisation (through national, regional and social media) and data collection coordination at national and/or regional levels. SFJN cleaned and analysed the data, and drafted the initial manuscript. All authors critically reviewed the draft and approved the final manuscript.

Funding

The study was supported by a grant from the European Research Council (ERC 671055) and VLIRUOS (the Flemish University cooperation). The study sponsors had no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the manuscript; and in the decision to submit the paper for publication.

Availability of data and materials

All the data presented in this article are available upon reasonable request from the ICPCovid consortium: [icpcovid@uantwerpen.be](https://icpcovid.uantwerpen.be)

Declarations

Ethics approval and consent to participate

This study was approved by the National Ethics Committee of Cameroon (Ref: 2020/05/1229/CE/CNERSH/SP of 06th May 2020) as well as the Ethics Committee of the University of Antwerp, Belgium (Ref: 20/13/148). All procedures were in accordance with international research guidelines. Only data from participants aged at least 18 years and who provided an informed e-consent were retained for analysis. All data were collected anonymously and treated with absolute confidentiality.

Consent for publication

Not applicable.

Competing interests

The authors have no conflicts of interest to declare.

Author details

¹Global Health Institute, University of Antwerp, Antwerp, Belgium. ²Brain Research Africa Initiative (BRAIN), Yaoundé, Cameroon. ³Brain Research Africa Initiative (BRAIN), Geneva, Switzerland. ⁴Division of Health Operations Research, Ministry of Public Health, Yaoundé, Cameroon. ⁵CDC Africa, African Union, Addis Ababa, Ethiopia.

Received: 28 January 2021 Accepted: 24 May 2021

Published online: 15 July 2021

References

- WHO Coronavirus Disease (COVID-19) Dashboard [Internet]. [cited 2021 Apr 1]. Available from: <https://covid19.who.int>

- Salari N, Hosseini-Far A, Jalali R, Vaisi-Raygani A, Rasoulopoor S, Mohammadi M, et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Glob Health*. 2020;16(1):57. <https://doi.org/10.1186/s12992-020-00589-w>.
- Kapila AK, Ram A, D'Uva L, Farid Y, Talib M. The importance of assessing the self-reported impact of COVID-19 on clinician mental health. *Eur J Psychiatry*. 2020;34(4):223–4. <https://doi.org/10.1016/j.ejpsy.2020.06.002>.
- Keubo FRN, Mboua PC, Tadongfack TD, Tchhoffo EF, Tatang CT, Zeuna JI, et al. Psychological distress among health care professionals of the three COVID-19 most affected regions in Cameroon: prevalence and associated factors. *Ann Med Psychol (Paris)*. 2021;179(2):141–6. <https://doi.org/10.1016/j.jamp.2020.08.012>.
- Ying Y, Ruan L, Kong F, Zhu B, Ji Y, Lou Z. Mental health status among family members of health care workers in Ningbo, China, during the coronavirus disease 2019 (COVID-19) outbreak: a cross-sectional study. *BMC Psychiatry*. 2020;20(1):379. <https://doi.org/10.1186/s12888-020-02784-w>.
- Pan X, Xiao Y, Ren D, Xu Z, Zhang Q, Yang L, et al. Prevalence of mental health problems and associated risk factors among military healthcare workers in specialized COVID-19 hospitals in Wuhan, China: A cross-sectional survey. *Asia-Pac Psychiatry*. 2020; [cited 2020 Nov 17]; Available from: <https://onlinelibrary.wiley.com/doi/10.1111/appy.12427>.
- Taquet M, Luciano S, Geddes JR, Harrison PJ. Bidirectional associations between COVID-19 and psychiatric disorder: retrospective cohort studies of 62 354 COVID-19 cases in the USA. *Lancet Psychiatry*. 2021;8(2):130–40. [https://doi.org/10.1016/S2215-0366\(20\)30462-4](https://doi.org/10.1016/S2215-0366(20)30462-4).
- Losada-Baltar A, Martínez-Huertas JA, Jiménez-Gonzalo L, del SM P-C, Gallego-Alberto L, Fernandes-Pires J, et al. Longitudinal Correlates of Loneliness and Psychological Distress During the Lockdown Situation due to COVID-19. Effects of Age and Self-Perceptions of Aging. *J Gerontol Series B*. 2021;gbab012.
- González-Sanguino C, Ausín B, Castellanos MÁ, Saiz J, López-Gómez A, Ugidos C, et al. Mental health consequences during the initial stage of the 2020 coronavirus pandemic (COVID-19) in Spain. *Brain Behav Immun*. 2020; 87:172–6. <https://doi.org/10.1016/j.bbi.2020.05.040>.
- AKMI B, Sakib N, Pakpour AH, Griffiths MD, Mamun MA. COVID-19-Related Suicides in Bangladesh Due to Lockdown and Economic Factors: Case Study Evidence from Media Reports. *Int J Ment Health Addict*. 2020; [cited 2020 Nov 17]; Available from: <http://link.springer.com/10.1007/s11469-020-00307-y>.
- Cameroon: WHO Coronavirus Disease (COVID-19) Dashboard [Internet]. [cited 2021 Jan 2]. Available from: <https://covid19.who.int>
- Government response strategy to the Coronavirus pandemic (COVID-19) | Prime Minister's Office [Internet]. [cited 2020 Nov 1]. Available from: <https://www.spm.gov.cm/site/?q=en/content/government-response-strategy-coronavirus-pandemic-covid-19>
- Mviena JLM, Fanne M, Gondo R, Mwamelo AJ, Esso L, Epée E, et al. How mental health care is changing in Cameroon because of the COVID-19 pandemic. *Lancet Psychiatry*. 2020;7(10):e62–3. [https://doi.org/10.1016/S2215-0366\(20\)30390-4](https://doi.org/10.1016/S2215-0366(20)30390-4).
- Cameroon Population (2020) - Worldometer. [cited 2020 Nov 17]. Available from: <https://www.worldometers.info/world-population/cameroon-population/>
- ICPCovid research group. International Citizen Project Covid-19. 2020 [cited 2020 May 11]. Available from: <https://www.icpcovid.com/>
- Ahorsu DK, Lin C-Y, Imani V, Saffari M, Griffiths MD, Pakpour AH. The Fear of COVID-19 Scale: Development and Initial Validation. *Int J Ment Health Addict*. 2020; [cited 2020 Apr 16]; Available from: <http://link.springer.com/10.1007/s11469-020-00270-8>.
- Parlapani E, Holvea V, Voitsidis P, Blekas A, Gliatas I, Porfyri GN, et al. Psychological and behavioral responses to the COVID-19 pandemic in Greece. *Front Psychiatry*. 2020;11:821. <https://doi.org/10.3389/fpsy.2020.00821>.
- Kroenke K, Spitzer RL, Williams JBW. The patient health Questionnaire-2: validity of a two-item depression screener. *Med Care*. 2003;41(11):1284–92. <https://doi.org/10.1097/01.MLR.0000093487.78664.3C>.
- Pence BW, Gaynes BN, Atashili J, O'Donnell JK, Tayong G, Kats D, et al. Validity of an interviewer-administered patient health questionnaire-9 to screen for depression in HIV-infected patients in Cameroon. *J Affect Disord*. 2012;143(1–3):208–13. <https://doi.org/10.1016/j.jad.2012.05.056>.
- Ueda M, Stickley A, Sueki H, Matsubayashi T. Mental health status of the general population in Japan during the COVID-19 pandemic. *Psychiatry Clin Neurosci*. 2020;74(9):505–6. <https://doi.org/10.1111/pcn.13105>.

21. Sigdel A, Bista A, Bhattarai N, Pun BC, Giri G, Marqusee H, et al. Depression, anxiety and depression-anxiety comorbidity amid COVID-19 pandemic: an online survey conducted during lockdown in Nepal: Public and Global Health; 2020. Available from: <http://medrxiv.org/lookup/doi/10.1101/2020.04.30.20086926>
22. Choi EPH, Hui BPH, Wan EYF. Depression and anxiety in Hong Kong during COVID-19. *Int J Environ Res Public Health*. 2020;17(10):3740. <https://doi.org/10.3390/ijerph17103740>.
23. Kanmogne GD, Qiu F, Ntone FE, Fonsah JY, Njamnshi DM, Kuate CT, et al. Depressive symptoms in HIV-infected and seronegative control subjects in Cameroon: Effect of age, education and gender. *PLOS ONE*. 2017;12: e0171956.
24. Ettman CK, Abdalla SM, Cohen GH, Sampson L, Vivier PM, Galea S. Prevalence of depression symptoms in US adults before and during the COVID-19 pandemic. *JAMA Netw Open*. 2020;3(9):e2019686. <https://doi.org/10.1001/jamanetworkopen.2020.19686>.
25. Lim GY, Tam WW, Lu Y, Ho CS, Zhang MW, Ho RC. Prevalence of depression in the community from 30 countries between 1994 and 2014. *Sci Rep*. 2018; 8(1):2861. <https://doi.org/10.1038/s41598-018-21243-x>.
26. Kannampallil TG, Goss CW, Evanoff BA, Strickland JR, McAlister RP, Duncan J. Exposure to COVID-19 patients increases physician trainee stress and burnout. *PLOS ONE*. 2020;15:e0237301.
27. Khanal P, Devkota N, Dahal M, Paudel K, Joshi D. Mental health impacts among health workers during COVID-19 in a low resource setting: a cross-sectional survey from Nepal. *Glob Health*. 2020;16(1):89. <https://doi.org/10.1186/s12992-020-00621-z>.
28. Vizheh M, Qorbani M, Arzaghi SM, Muhidin S, Javanmard Z, Esmaili M. The mental health of healthcare workers in the COVID-19 pandemic: a systematic review. *J Diabetes Metab Disord*. 2020;26;19(2):1-12. <https://doi.org/10.1007/s40200-020-00643-9>.
29. Pearman A, Hughes ML, Smith EL, Neupert SD. Mental health challenges of United States healthcare professionals during COVID-19. *Front Psychol*. 2020;11:2065. <https://doi.org/10.3389/fpsyg.2020.02065>.
30. Agberotimi SF, Akinsola OS, Oguntayo R, Olaseini AO. Interactions between socioeconomic status and mental health outcomes in the Nigerian context amid COVID-19 pandemic: a comparative study. *Front Psychol*. 2020;11: 559819. <https://doi.org/10.3389/fpsyg.2020.559819>.
31. Mboua CP, Keubo FRN, Fouaka SGN. Anxiété et dépression associées à la prise en charge de la COVID-19 chez les personnels de santé au Cameroun. *Evol Psychiatr (Paris)*. 2021;86(1):131–9.
32. Hawryluck L, Gold WL, Robinson S, Pogorski S, Galea S, Styra R. SARS control and psychological effects of quarantine, Toronto, Canada. *Emerg Infect Dis*. 2004;10(7):1206–12. <https://doi.org/10.3201/eid1007.030703>.
33. Passos L, Prazeres F, Teixeira A, Martins C. Impact on mental health due to COVID-19 pandemic: cross-sectional study in Portugal and Brazil. *Int J Environ Res Public Health*. 2020;17(18):6794. <https://doi.org/10.3390/ijerph17186794>.
34. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395(10227):912–20. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8).
35. Reicher S, Drury J. Pandemic fatigue? How adherence to covid-19 regulations has been misrepresented and why it matters. *BMJ*. 2021; [cited 2021 Apr 27]. Available from: <https://blogs.bmj.com/bmj/2021/01/07/pandemic-fatigue-how-adherence-to-covid-19-regulations-has-been-misrepresented-and-why-it-matters/>.
36. Fouogue JT, Noubom M, Kenfack B, Dongmo NT, Tabou M, Megozeu L, et al. Poor knowledge of COVID-19 and unfavourable perception of the response to the pandemic by healthcare workers at the Bafoussam Regional Hospital (West Region-Cameroon). *Pan Afr Med J*. 2020;37. [cited 2020 Nov 12] Available from: <https://www.panafrican-med-journal.com/content/series/37/1/1/full>
37. Yao H. The more exposure to media information about COVID-19, the more distressed you will feel. *Brain Behav Immun*. 2020;87:167–9. <https://doi.org/10.1016/j.bbi.2020.05.031>.
38. COVID-19 Map. Johns Hopkins Coronavirus Resour. Cent. [cited 2020 Apr 15]. Available from: <https://coronavirus.jhu.edu/map.html>
39. Xiang Y-T, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry*. 2020;7(3):228–9. [https://doi.org/10.1016/S2215-0366\(20\)0046-8](https://doi.org/10.1016/S2215-0366(20)0046-8).
40. Bakehe NP, Fambeu AH, Tamokwe Paptie GB. Internet adoption and use in Cameroon. Cameroon: Theoretical and Applied Economics Research Group, University of Douala; 2017.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions



Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com